

## Chapter 8

### Vapor Pressure

#### Solutions

##### Problem 8.1\* *Basic Vapor Pressure Calculations*

(a) Prepare the data to be expressed as in the linear relationship Eq. 8.7. Note that at 45.0 and 74.6°C, TeMB is a solid. Hence, group the data according to the aggregate state of the compound.

Perform least squares fits of  $\ln p_i^*$  versus  $1/T$  (see Figure below). The results are:

$$\text{Solid compound: } \ln p_{is}^* (\text{mm Hg}) = -\frac{8609\text{K}}{T} + 27.1 \quad (1)$$

$$\text{Liquid compound: } \ln p_{iL}^* (\text{mm Hg}) = -\frac{5676\text{K}}{T} + 18.7 \quad (2)$$

If you would have converted mm Hg to bar (1 mm Hg = 0.01333 bar), the intercepts of Eqs. 1 and 2 would be 20.5 and 12.1, respectively.

Insert  $T = 293.2$  K (20°C) into Eq. 1, calculate  $p_{is}^*$ :

$$p_{is}^* (20^\circ\text{C}) = 0.10 \text{ mm Hg} = 0.000133 \text{ bar} = 13.3 \text{ Pa}$$

Insert  $T = 423.2$  K (150°C) into Eq. 2, calculate  $p_{iL}^*$ :

$$p_{iL}^* (150^\circ\text{C}) = 198 \text{ mm Hg} = 0.264 \text{ bar} = 26400 \text{ Pa}$$

Hence, the vapor pressure of TeMB is more than three orders of magnitude greater at 150°C than at 20°C, illustrating the strong temperature dependence of this compound property.

(b) Insert  $T = 293.2$  K into Eq. 2 to estimate the vapor pressure of subcooled TeMB at 20°C:

$$p_{iL}^* (20^\circ\text{C}) = 0.52 \text{ mm Hg} = 0.00069 \text{ bar} = 69 \text{ Pa}$$

Hence, at 20°C,  $p_{iL}^*$  is about five times larger than  $p_{is}^*$ . Remember, however, that you have extrapolated this value over quite a large temperature range. Use Eq. 8-9 to calculate  $\Delta_{\text{fus}}G_i$ :

$$\Delta_{\text{fus}}G_i (20^\circ\text{C}) = RT \ln \frac{p_{iL}^*}{p_{is}^*} = (2.44 \text{ kJ mol}^{-1}) \ln \frac{(0.00069)}{(0.000133)} = 4.0 \text{ kJ mol}^{-1}$$


Estimate  $\Delta_{\text{fus}}H_i$  from the average  $\Delta_{\text{vap}}H_i$  and  $\Delta_{\text{sub}}H_i$  that you can derive from the slopes of the regression lines Eqs. 1 and 2 (see also Eqs. 8-10 and 8-11):

$$\Delta_{\text{fus}}H_i = \Delta_{\text{sub}}H_i - \Delta_{\text{12}}H_i = R \cdot \text{slope 1} - R \cdot \text{slope 2} = 24.4 \text{ kJ mol}^{-1},$$

and since  $\Delta_{\text{fus}}S_i = (\Delta_{\text{fus}}H_i - \Delta_{\text{fus}}G_i) / T$ , you get

$$\Delta_{\text{fus}}S_i = 69.6 \text{ J mol}^{-1}\text{K}^{-1}$$

Note again that in these calculations, all  $\Delta_{12}H_i$  values have been assumed to be constant over the temperature range considered. Therefore, all changes in  $\Delta_{\text{fus}}G_i$  (which is zero at  $T_m$ ) are attributed to a change in  $\Delta_{\text{fus}}S_i$  with changing temperature. This is, of course, not exactly correct.



Temperature dependence of the vapor pressure of TeMB: Plot of  $\ln p_i^*$  versus  $1/T$ .

### Problem 8.2 A Solvent Spill in Your Class Room

In the temperature range considered, PCE is a liquid. Hence, all four data points can be used to establish the vapor pressure-temperature equation. In Appendix C (Wiley website) you find the molecular weight ( $165.83 \text{ g mol}^{-1}$ ), the density at  $20^\circ\text{C}$  ( $1.62 \text{ g cm}^{-3}$ ), and the Abraham parameters ( $V_i=0.84$ ,  $L_i=3.58$ ,  $S_i=0.44$ ,  $A_i=0$ ,  $B_i=0$ ) of PCE. At  $20^\circ\text{C}$  10 L of PCE correspond, therefore, to 16.2 kg or 97.69 moles.

(a) The linear regression of the four data points ( $\ln p_i^*$  vs.  $1/T$ ) yields:

$$\ln p_i^* (\text{kPa}) = -4609 (1/T) + 16.36$$

Insertion of  $T = 293\text{K}$  gives a  $p_i^*$  value at  $20^\circ\text{C}$  of 1.88 kPa or a maximum PCE concentration ( $p_i^*/RT$ ) =  $(0.0188\text{bar}/24.35 \text{ L bar mol}^{-1}) = 0.77 \text{ mmol L}^{-1}$  or  $128 \text{ g m}^{-3}$ . This is more than 4 orders of magnitude above the odor threshold. The total amount of PCE evaporated in the  $50 \text{ m}^3$  is then about 6.4 kg (4 L) or about 40% of the PCE that was originally in the bottle.

Using Eq. 8-19 yields a vapor pressure of about 1.7 kPa at  $25^\circ\text{C}$ . With Eq. 8-14, one may use this estimated vapor pressure to estimate PCE's enthalpy of vaporization to be  $42 \text{ kJ mol}^{-1}$ . With this enthalpy and the vapor pressure at 298K, Eq. 8-7 can be used to calculate the vapor pressure at 293K as 1.3 kPa. This estimate corresponds quite well with the value extrapolated from the experimental data (much less than a factor of 2).

(b) At  $80^\circ\text{C}$  all PCE has evaporated yielding a maximum concentration of  $16.2 \text{ kg} / 15 \text{ m}^3 = 1.08 \text{ kg m}^{-3}$  which is about 5 orders of magnitude above the odor threshold!

### Problem 8.3 *How Much Freon is Left in the Old Pressure Bottle Left at a Dump Site?*

At 0°C the minimum pressure required to have liquid Freon 12 in the bottle is more than 300 kPa or 3 bar, hence at 10°C the pressure would have to be higher. Therefore you can assume that there is no liquid Freon 12 present in the bottle, and you can estimate the amount present:

$C_{\text{bottle}} = p_i/RT = 2.7 \text{ bar}/(23.54 \text{ bar L mol}^{-1}) = 0.115 \text{ mol L}^{-1}$ ; this yields a total amount of 0.34 mol or 41.6 g Freon 12 left in the bottle.

The free energy of condensation at 25°C is given by (Eq. 8.2):

$$\Delta_{\text{cond}}G_i = RT \ln p_i^* = (2.48 \text{ kJ mol}^{-1})(\ln 6.51) = 4.6 \text{ kJ mol}^{-1}$$

To get the enthalpy of condensation, calculate the linear regression of the five data points ( $\ln p_i^*$  vs.  $1/T$ ):

$$\ln p_i^* (\text{kPa}) = -2437 (1/T) + 14.65$$

Since  $\Delta_{\text{cond}}H_i = (\text{slope})R$ , you get an enthalpy of condensation of  $-20.3 \text{ kJ mol}^{-1}$ , and an entropy of condensation,  $\Delta_{\text{cond}}S_i = (\Delta_{\text{cond}}H_i - \Delta_{\text{cond}}G_i)/T = (-20.3 \text{ kJ mol}^{-1} - 4.6 \text{ kJ mol}^{-1})/298\text{K}$  of  $-83.6 \text{ J mol}^{-1}\text{K}^{-1}$ .

**Problem 8.4 What Are the Differences Between Freon 12 And its Replacement HFC-134a? (From Roberts, 1995)**

The replacement was necessary because, due to its large residence time in the atmosphere, Freon 12 is both an “ozone killer” as well as a very potent greenhouse gas. HFC-134a does not react with stratospheric ozone and has a shorter residence time due to reaction with OH-radicals. However, it is still a potent greenhouse gas.

(a) The normal boiling point,  $T_b$ , is the temperature at which the vapor pressure is 1 bar or 100kPa (Section 8). From the vapor pressure temperature relationship derived for Freon 12 in Problem 8.3:

$$\ln p_i^* (\text{kPa}) = -2437 (1/T) + 14.65$$

you obtain a boiling point of 242.8 K or  $-30.4^\circ\text{C}$  for this compound. The relationship derived for HFC-134a is:

$$\ln p_i^* (\text{kPa}) = -2748 (1/T) + 15.75$$

which yields a  $T_b$  value of 246.7 K or  $-26.5^\circ\text{C}$ . Note that above the boiling point, the vapor pressure reflects the vapor pressure of the superheated liquid (see Fig. 8.2).

(b) Setting the two equations equal and solving for  $T$  yields a  $T_{\text{equal}}$  of 283K or about  $10^\circ\text{C}$ .

(c) As calculated in Problem 8.3, for Freon 12,  $\Delta_{\text{vap}}H_i = 20.3 \text{ kJ mol}^{-1}$  and  $\Delta_{\text{vap}}S_i = 83.6 \text{ J mol}^{-1} \text{ K}^{-1}$  at  $25^\circ\text{C}$ ). The same calculation for HFC-134a yields  $\Delta_{\text{vap}}G_i = -5.5 \text{ kJ mol}^{-1}$ , and thus  $\Delta_{\text{vap}}H_i = 22.8 \text{ kJ mol}^{-1}$  and  $\Delta_{\text{vap}}S_i = 94.0 \text{ J mol}^{-1} \text{ K}^{-1}$  for HFC-134a.

(d) Insertion of  $T = 323.2\text{K}$  into the above equations yields the following vapor pressures at  $50^\circ\text{C}$ :

$$\text{Freon 12: } p_i^* = 12.1 \text{ bar} \quad ; \quad \text{HFC-134a: } p_i^* = 13.9 \text{ bar}$$

Hence the difference is more than 10%, and some adjustments had to be made in the air conditioners in cars when Freon 12 was replaced by HFC-134a.

### Problem 8.5 A Public Toilet Problem

In Appendix C (Wiley website) you find the molecular weight ( $147.0 \text{ g mol}^{-1}$ ) and the Abraham parameters ( $V_i=0.96$ ,  $L_i=4.44$ ,  $S_i=0.75$ ,  $A_i=0$ ,  $B_i=0$ ) of 1,4-DCB.

(a) At  $20^\circ\text{C}$  1,4-DCB is a solid. Hence you can only use the first four data points for deriving the vapor pressure – temperature relationship:

$$\ln p_{is}^* (\text{kPa}) = -5458(1/T) + 17.08$$

Insertion of  $T = 293\text{K}$  yields  $p_{is}^* = 0.21 \text{ kPa}$  or  $2.1 \times 10^{-3} \text{ bar}$ , which corresponds to a maximum concentration of  $p_{is}^*/RT = 8.7 \times 10^{-5} \text{ mol L}^{-1}$  or about  $13 \text{ g m}^{-3}$ .

(b) Inserting  $T_b$  into Eq. 8-17 (with  $K_F=1$ ; since dipole moment is zero for this compound, Fishtine, 1963) yields the vapor pressure of the subcooled liquid 1,4-DCB at  $20^\circ\text{C}$ :  $\log p_{iL}^* (\text{bar}) = -2.8$  or  $p_{iL}^* = 1.5 \times 10^{-3} \text{ bar}$ . Using Eq. 8-21 with  $\Delta_{\text{fus}} S_i(T_m) = 45 \text{ J mol}^{-1}\text{K}$  (Table 8.2) the yields a solid vapor pressure of  $0.87 \times 10^{-3} \text{ bar}$  or  $5.3 \text{ g m}^{-3}$ .

(c) Using the pp-LFER Eqs. 8-19 and 8-14 yields a subcooled liquid vapor pressure at  $20^\circ\text{C}$  of  $145 \text{ Pa}$  or  $1.45 \times 10^{-3} \text{ bar}$ , which is virtually the same value as predicted from the boiling point (b).

Hence, the predictions are about a factor of two smaller than the value obtained from the experimental data which is quite satisfactory.

In the Death Valley, 1,4-DCB is a liquid. Using the boiling point ( $100 \text{ kPa}$ ) and the vapor pressure at  $99^\circ\text{C}$  ( $10 \text{ kPa}$ ) given in the Table above, one obtains

$$\ln p_{iL}^* (\text{kPa}) = -5132(1/T) + 16.09$$

and thus  $p_{iL}^* (\text{kPa})$  at  $60^\circ\text{C} = 1.97 \text{ kPa}$  or  $1.99 \times 10^{-2} \text{ bar}$  or about  $105 \text{ g m}^{-3}$  which is an eight times higher concentration than at  $20^\circ\text{C}$ , demonstrating the strong temperature dependence of the vapor pressure of the compound.

### Problem 8.6 True or False?

In Appendix C you find

1,2-DCB:  $T_m = -17^\circ\text{C}$ ,  $T_b = 180^\circ\text{C}$ ; 1,4-DCB:  $T_m = 53.1^\circ\text{C}$ ,  $T_b = 174^\circ\text{C}$ . Hence, at  $60^\circ\text{C}$  both compounds are liquids, and you would expect that the one with the lower boiling point (1,4-DCB) would exhibit the higher vapor pressure, which is of course, reflected in the corresponding estimated vapor pressures. However, at  $20^\circ\text{C}$ , this is not obvious, since one has to take into account that, in contrast to 1,2-DCB, 1,4-DCB is a solid at this temperature.

For estimating the (subcooled) liquid vapor pressures, insert the respective  $T_b$  values (in K) into Eq. 8-17. Note, according to Fishtine (1963), for substituted benzenes the  $K_F = 1 + 2\mu/100$  where  $\mu$  is a compound's dipole moment. Since the chlorine substituents on 1,4-DCB are symmetrically opposed to one another,  $\mu$  for 1,4-DCB is 0 and  $K_F = 1$ . But 1,2-DCB has permanent dipole moment of 2.5 (Haynes, 2014), so it's  $K_F$  is 1.05). The resulting  $p_{iL}^*$  values are

at  $60^\circ\text{C}$ :

1,2-DCB :  $1.2 \times 10^{-2}$  bar

1,4-DCB :  $1.8 \times 10^{-2}$  bar

at  $20^\circ\text{C}$ :

1,2-DCB :  $0.91 \times 10^{-3}$  bar

1,4-DCB :  $1.7 \times 10^{-3}$  bar

However correcting the vapor pressure of the 1,4-DCB solid (Eq. 8-21 with  $\Delta_{\text{fus}} S_i(T_m) = 45 \text{ J mol}^{-1}\text{K}$ ;

Table 8.2) yields the solid (real) vapor pressure of  $0.92 \times 10^{-3}$  bar.

The person is slightly wrong to the extent one can trust the estimates!

For determining at which temperature the two isomers exhibit the same vapor pressures, you need to have the vapor pressure – temperature relationship for the two compounds over the temperature range of the intersect of the two lines which has to be between  $20^\circ\text{C}$  and the melting point temperature of 1,4-DCB. For 1,2-DCB you can take the vapor pressures at  $20^\circ\text{C}$  and  $60^\circ\text{C}$  that you have already calculated. For 1,4-DCB you need to calculate the vapor pressure at the melting point, hence at  $53.1^\circ\text{C}$  (Eq. 8-17). The value is  $1.28 \times 10^{-2}$  bar. This yields the following vapor pressure-temperature relationships:

for 1,4-DCB:

$$\ln p_{is}^* (\text{bar}) = -7617(1/T) + 19.00$$

and for 1,2-DCB:

$$\ln p_{iL}^* (\text{bar}) = -6245(1/T) + 14.31$$

setting the two equations equal yields the temperature at which the vapor pressures are equal.

$$T = 292.5\text{K or } 19.5^\circ\text{C}$$

### Problem 8.7 Vapor pressure of Alkyl Nitriles from Biomass Burning

The melting temperature and boiling point indicate that dodecyl nitrile (DN) is a liquid at 25°C. So one may use Eq. 8-17 to estimate this compound's vapor pressure:

$$\log p_{\text{DNL}}^* (\text{bar}) = -(1/2.303)K_{\text{F}}(4.4 + \ln T_{\text{b}}) \left[ 1.8 \left( \frac{T_{\text{b}}}{T} - 1 \right) - 0.8 \ln \frac{T_{\text{b}}}{T} \right]$$

where  $K_{\text{F}}$  is DN's "Fishtine" factor. For nitriles with long (>12C) alkyl chains like DN, the  $K_{\text{F}}$  value is 1.01 (Fishtine, 1963). Hence, one may use this small factor and DN's boiling temperature (277°C = 550 K) to estimate its vapor pressure:

$$\log p_{\text{DNL}}^* (25^\circ\text{C}, \text{bar}) = -(1/2.303)(1.01)(4.4 + \ln 550) \left[ 1.8 \left( \frac{550}{298} - 1 \right) - 0.8 \ln \frac{550}{298} \right] = 1.4 \times 10^{-5} \text{ bar} = 1.4 \text{ Pa}$$